

TOWARDS QUANTIFYING NON-LOCAL INFORMATION TRANSFER: FINITE-BIT NON-LOCALITY¹

Michael Steiner

mjs@mike.nrl.navy.mil

February 3, 1999

Naval Research Laboratory, Code 5340, Washington, DC 20375

Abstract

The advent of Bell's inequalities provoked the possibility that entangled quantum phenomena is non-local in nature. Since teleportation only requires a finite amount of classical information, i.e. two bits, the author asks whether or not it is possible to further characterize the nature of internal correlation in terms of information. Towards this end, the issue of the amount of information that is transferred internally and non-locally is addressed. There are two possibilities: the amount is infinite or the amount is finite. A partial answer to this problem is given: it is shown that models exist whereby the amount is finite. The EPR-Bell cosine correlation can be reproduced exactly using on average 1.48 bits. The issue of simultaneity and the problems it poses are also examined in this context. Several extensions are suggested.

I. BACKGROUND

Since the advent of Bell's inequalities and the subsequent verification of the of the Quantum Mechanical predictions, it is speculated by some that there exists in nature information transfer across large distances that occurs faster than the speed of light, and perhaps, instantaneously. While such information transfer may occur internally to elementary particles, it is widely accepted that internal information transfer cannot be utilized to develop a classical signaling system that transmits external information instantaneously. However, that such

¹Paper presented at the Physics of Quantum Electronics Conference, Jan. 3-7, 1999

internal information transfer may occur in nature is counter-intuitive. The purpose of this paper is to develop a framework towards determining the amount of internal information (nonredundant) that is internally transferred non-locally.

The author is motivated to investigate this for two reasons. Firstly, it is seen in the teleportation experiments that in addition to the Einstein-Podolsky-Rosen (EPR) correlation, only a finite amount of classical information is required, yet a particle with an arbitrary density matrix is teleported. Hence, it is worthwhile to see if the teleported particle can actually be specified with only the initial state of the EPR particles, and finite-bit non-locality (FBNL). Certainly, if only infinite-bit NL models could be proven to exist and no FBNL model exists, then the specification of a particle at a distance would require an infinite amount of information, and the fact that teleportation is possible with only two classical bits would be astonishing from an information perspective. If it can be proven that the amount of non-locality required to replicate the teleportation experiments is greater than 2 bits, then teleportation must be utilizing non-locality (considered over the set of FBNL models). If the amount of non-locality required is less than or equal to 2 bits, then a model may exist whereby the 2 bits that are classically transmitted contain the information of the state of the particle, and the initial local hidden information is simply used as a reference. Related questions have been asked by others, for example, Vaidman in [1] questions if the essence of a state of a spin-1/2 particle is just 2 bits.

The second reason to investigate whether or not a FBNL model exists is that an infinite amount of information can only be transferred via a channel that is effectively noiseless. That is, since the capacity of a typical channel behaves like $B \log(1 + \text{SNR})$, where B is the bandwidth and SNR denotes the signal-to-noise ratio, the only way such information could be exchanged is if the SNR is infinite ($B = \infty$ does not work). Hence, if no FBNL models exists, then one is forced to accept that the internal channel is effectively noiseless. This is hard to accept, since no analogy exists classically. On the other hand, no classical analogy exists for many quantum phenomenon, hence this second reason is not as pervasive as the first.

In Section II, non-local information transfer will be formally defined. In Section III, we will show several admissible models, including the finite-bit non-locality model. It is known that models of non-locality such as Bohm's model as well as the models considered in this paper have not been reconciled with special relativity and in particular, the issue of simultaneity. In Section IV implications of this issue are examined

II. NON-LOCAL INTERNAL INFORMATION TRANSFER

There have been numerous studies regarding quantum information. However, the problem presented in this paper does not appear to have been solved (related results have recently been obtained independently by Brassard, Cleve, and Tapp [2] discussed in Section V). In [3] the question of how many bits are required to transmit a qubit reliably is examined. The approach appears to be based on an approximation, and a fidelity or accuracy is defined. In [4] several related information quantities are addressed.

In order to understand entanglement it is useful (in my opinion) to consider models. The class of models assumed here is that a positive amount of hidden information is transferred between particles. The information is hidden in the sense that it cannot be manipulated by external observers for the purpose of sending superluminal signals. In this paper, it is shown that a finite amount of information can be used to quantify the amount of hidden information. In fact, a technique is shown whereby only 1.48 bits is required to replicate exactly the cosine correlation function expected for the EPR experiment (with spin measurement).

A model that provides the same expected values as quantum mechanics for a given experiment will be termed an admissible model for the experiment. The reader may note that the wavefunction in quantum mechanics itself is an admissible model. However, it could be argued that the wavefunction in quantum mechanics is only a partially complete model. Additionally, it is well known that the measurement problem, or wavefunction collapse in the Copenhagen interpretation of quantum mechanics, does not have a well-defined model [5].

The issue of how much, if any, information must be transferred between particles in order to observe the correlations observed in the EPR experiment deserves attention. There are three possibilities for a given reference frame: 1) no information is transferred, 2) a finite amount is transferred, and 3) an infinite amount of information is transferred. A reference frame is assumed whereby the measurement devices are not moving and the EPR source is fixed closer to one measurement device than the other. The problem of Lorentz invariance and the issue of simultaneity is examined in Section IV.

Although the first possibility above whereby no information is transferred is discussed at the end of Section III, the primary purpose of the paper is to investigate whether or not it is possible to model the particles and obtain the quantum mechanical correlation with FBNL as opposed to infinite information models.

III. INTERNAL INFORMATION TRANSFER MODELS

The experimentally shown violation of Bell's inequalities points to the possible existence of non-local phenomenon in nature. However, neither Bell's inequality nor quantum mechanics shed much light on just how these non-local effects propagate. Bohm's and other theories do provide some insight, but they require that some non-local effects occur. For example, the quantum potential can change instantaneously in Bohm's theory. However, it appears from the equations that specify the trajectories for this experiment [6][p. 132] that the trajectories are a function of the parameters of both Stern-Gerlach devices. Hence Bohm's theory does not shed light on how much non-locality is required.

We will examine the amount of information that is required to be transferred via models that allow for the non-local transfer of hidden information. Consider the EPR experiment shown in Fig. 1. There is an initial state Λ_I that characterizes the initial state. Experimenter A has a Stern-Gerlach measurement apparatus oriented at θ_a which records $A = 1$ or -1 and experimenter B has a similar apparatus specified by θ_b . The source is positioned closer to A than B. Two entangled spin $1/2$ particles will be assumed to be emitted with state $|\psi\rangle = \frac{1}{\sqrt{2}}(|\uparrow\rangle|\downarrow\rangle - |\downarrow\rangle|\uparrow\rangle)$

We will now assume the possibility that classical hidden information denoted by Λ_H can be transferred through a channel with capacity C_H bits/use, and that this channel is used once so that only C_H bits can be transferred. We assume that Λ_H is chosen from a discrete memoryless source so that its information content may be finite, given by $H(\Lambda_H)$. Note that specifying a continuous value exactly requires infinite precision. In more precise terms, trying to send a continuous source through a channel requires a capacity specified by the rate distortion function $R(d)$ in information theory. As long as one allows for a given amount of distortion, the information can be sent reliably. However, if no distortion is allowed, i.e. $d = 0$, then for most continuous sources (such as the Gaussian source) $R(d)$ is unbounded. Since we do not require to transmit the set Λ_I at a distance, we assume that Λ_I is a set of fixed complex numbers. However, whether or not this assumption is required will be examined in Section V.

Denote a finite-bit model (FBM) $M = \{\Lambda, R\}$ to be a set of variables $\Lambda = \lambda_I, \lambda_H(\theta_A)$ and a measurement rule $R = \{A(\lambda_I, \theta_a), B(\lambda_I, \theta_b, \lambda_H)\}$. A model that agrees statistically with quantum mechanics for a given experiment $\mathcal{E}$ will be called an admissible model for that experiment and is denoted by $M \in \mathcal{A}(\mathcal{E})$. Finite bit admissible models are only a

subset of $\mathcal{A}(\mathcal{E})$ and the admissible models for this class are denoted $\mathcal{A}_{\text{FB}}(\mathcal{E})$. The quantum mechanical predictions for this experiment are

$$E(AB) = -\cos(\theta_a - \theta_b) \quad (1)$$

with marginals $E(A) = 0$ and similarly for B .

Now, there are two cases to be considered. The first is where C_H is an average constraint so that the amount of information can be more than C_H on any one channel use. The second is a peak constraint so that for each channel use, only C_H bits or less are transmitted. Models M for which there exists a source code characterizing Λ_H with lengths that are all less than or equal to C_H bits are called peak constrained models and are denoted as $\mathcal{M}_{\text{peak}}$. Similarly models for which the source code has average lengths less than C_H are denoted $\mathcal{M}_{\text{avg}}$. In this paper, only the average constraint will be examined. See Section V for results regarding the peak constraint. Define

$$C_{H,\text{avg}} = \min_{M \in \mathcal{A}_{\text{FB}}(\mathcal{E}), \mathcal{M} \in \mathcal{M}_{\text{avg}}} H(\Lambda_H). \quad (2)$$

and similarly for $C_{H,\text{peak}}$. Note that $C_{H,\text{avg}} \leq C_{H,\text{peak}}$. We will now demonstrate an average constrained model for which $H(\lambda_H) = 1.48$ and hence $C_{H,\text{avg}} \leq 1.48$ (for this experiment).

Generating an Arbitrary Variate at a Distance with Finite Information

Consider first the following more general problem. Suppose that one would like to generate at B a single random variate which has any distribution $f_{\theta_a}(x)$ that is parameterized by a continuous parameter such as θ_A . For example $f_{\theta_a}(x)$ could be a distribution with a peak at $x = a$. Suppose furthermore that both A and B have available to them a random number seed specified by Λ_I , and some method of generating random numbers which is composed of variates from some *a priori* distribution. However, only A knows θ_a . Is it possible for A to send on average a finite amount of information to B so that B can construct a single random variate with distribution exactly equal to $f_{\theta_a}(x)$? The answer is surprisingly yes, at least under certain conditions. One solution is afforded by adopting the rejection method of non-uniform random number generation[7]. The rejection method is as follows. Suppose that one can find a distribution $g(x)$ and a constant $c \geq 1$ so that

$$f_{\theta_a}(x) \leq cg(x)$$

for all x and all θ_a . Perform the following steps:

Step 1: Generate a variate w with distribution $g(x)$ and another variate u that is uniformly distributed on $[0, 1]$.

Step 2: Compute the function $T = u c g(w)/f(w)$.

Step 3: If $T > 1$ repeat starting from step 1. If $T \leq 1$ then w has the desired distribution $f_{\theta_a}(x)$.

Now, all that is needed for A to do is to send to B the iteration number where he first succeeded in getting $T \leq 1$. B then simply generates this variate using the same seed and has succeeded in generating a variate from the distribution f_{θ_a} . One should note that the amount of information required may or may not be finite since potentially the iteration at which A succeeds is not bounded by any constant. Hence, the convergence of this process are significant. It is known that the convergence of this process is only a function of the parameter c , and is specified by the geometric distribution, $P(K = k) = (1 - p)^{k-1}p$, where $p = 1/c$ and K is the iteration number. In fact $P(K > k) = (1 - p)^k \leq \exp(-pk)$, hence the technique converges exponentially fast. From source coding theory, it is known that a discrete memoryless source can be coded by a variable length code with average length equal (or nearly so) to its entropy. This is also the average number of bits that A must send to B. The entropy of this process is given by

$$H = \sum_k -P(K = k) \log P(K = k)$$

which can be shown to be

$$H = -\log p - \frac{(1 - p)}{p} \log(1 - p). \quad (3)$$

Hence if one can find a $c \geq 1$ whereby $cg(x)$ that is uniformly larger than $f_{\theta_a}(x)$, then a technique follows which only requires a finite amount of information.

Application to Bell pair

This technique can be immediately applied to the Bell pair. It was shown in [8] that if we define $A(x, \theta_a) = \text{sign}(\cos(x - \theta_a))$, $B(x, \theta_b) = -\text{sign}(\cos(x - \theta_b))$ and $P(x, \theta_a) = 1/4|\cos(x - \theta_a)|$ then,

$$-\cos(\theta_a - \theta_b) = \int A(x, \theta_a)B(x, \theta_b)P(x, \theta_a)dx \quad (4)$$

and $E(A) = E(B) = 0$. The rejection method above is applied at particle A to generate $f_{\theta_a}(x) = 1/4|\cos(x - \theta_a)|$ using a random number generator with seed specified by λ_I and with $g(x)$ uniformly distributed on $[0, 2\pi]$. We assume that a new seed is generated for every new experiment with a Bell pair. Note that the condition $f_{\theta_a}(x) \leq cg(x)$ is satisfied for $c = 2\pi/4$ for all θ_a . Particle A measures $\text{sign}(\cos(x - \theta_a))$ and sends to B the iteration number k where A first succeeded in $T \leq 1$. Particle B then generates x from seed λ_I and can now directly generate $B(x, \theta_b) = \text{sign}(\cos(x - \theta_b))$ and therefore has succeeded in exactly generating the cosine correlation function. One can check that all marginals and the joint distribution are in agreement with the Quantum Mechanical results as well. For this problem, the information required is specified by Eqn. (3) with $p = 1/c = \frac{4}{2\pi}$. In this case Eqn. (3) reduces to $H = 1.48$ bits. Hence for this experiment, $C_{H,avg} \leq 1.48$ bits. A simulation was written and both the cosine correlation function and average number of bits equal to 1.48 were verified.

Other models and a lower bound

Note that the FB models considered have a particular form: $M = \{\Lambda, R\}$, $\Lambda = \lambda_I, \lambda_H(\theta_A)$ and a measurement rule $R = \{A(\lambda_I, \theta_a), B(\lambda_I, \theta_b, \lambda_H)\}$. This constraint on the form of the model means that we are considering only the subset $\mathcal{A}_{\text{FB}}(\mathcal{E}) \subset \mathcal{A}(\mathcal{E})$. Note that upper bounds on a FB model class are still valid bounds if we enlarge the class of models. However, a non-zero lower bound on $C_{(H,\cdot)}$ obtained for the class of FB models may not be valid if we consider an enlarged model set. Hence $C_{(H,\cdot)} = 0$ can not be ruled out very easily. For example, a class of models could be envisioned that assumes that a deterministic function exists so that all future events can be predicted locally given an initial state. If such a model exists it is plausible that no non-locality is required to explain the EPR experiments. However, it is not clear that such a model exists. If we restrict our model to the FB class, then it appears that a lower bound should be attainable for at least this set.

IV. ISSUES OF LORENTZ INVARIANCE AND SIMULTANEITY

It has been noted in several papers such as [9] among others that realistic theories either require a preferred frame of reference or will give rise to well-known causal paradoxes. This has implications on causality when considering multi-simultaneity experiments [10]. There are several types of causality violations that can emerge when considering such models. We

will consider two types. The first type of violation is that the experimenter can observe what he is about to do. The second violation is that the experimenter can not determine what he is about to do, but what he is about to do must be some function of the future (but he can not determine what the function is). This second violation is less serious than the first type, but bothersome nonetheless. Interestingly, it turns out that the first type of violation can not occur with our model. This is due to the fact that the distribution of the information that is sent from particle A to B is a geometric distribution which is independent of the parameters of the detection devices, such as θ_a . Furthermore, since the seed is hidden information and is non-accessible, it is not possible for any external observers in any reference frames to glean anything about their future. However, it appears that multi-simultaneity experiments could be devised as in [10] that would show the second type of violation occurring.

Note that such problems are not limited to this theory. It is well known that other realistic models such as Bohm's theory, have problems with Lorentz invariance. However, this problem still appears to be significant. The relationship between entanglement and simultaneity has not been deeply explored. Some related work on this was done recently by Suarez [11]. This problem is also related to collapse models. Both non-relativistic and relativistic wave function collapse models are reviewed in [12].

V. FURTHER WORK

There is significant further work that can be done. Note that $C_{H,avg}$ was not determined for general polarization but for a particular polarization such as linear polarization. It would be useful to generalize this result both for general spin 1/2 particles and general measurements. In this paper, the EPR experiment with two particles was analyzed. However, it would be useful to analyze the teleportation problem which requires three particles in more detail. If it can be proven that the amount of non-locality required to replicate the teleportation experiments is greater than 2 bits, then teleportation must be utilizing non-locality in addition to the 2 bits (considered over the set of FBNL models). If, on the other hand, the amount of non-locality required is less than or equal to 2 bits, then a model may exist whereby the 2 bits that are classically transmitted contain the state of the particle (in addition to the shared information λ_I).

The result that a finite amount of information can be used to obtain the cosine correlation function was also obtained independently by Brassard, Cleve, and Tapp [2]. In fact, they show that 4 bits can be used to obtain the cosine correlation function, hence have found

an upper bound on $C_{H,peak}$. They also generalized their work to arbitrary von Neumann measurements and showed that 8 bits would suffice. This is very interesting, and their approach is fairly different than the rejection method considered here. Their main interest is in Quantum Simulation, which is an application that the rejection method could also be applied to. Hence we now know that $C_{H,peak} \leq 4$ and $C_{H,avg} \leq 1.48$ (for generation of the cosine correlation function).

We have examined FB non-locality, but with λ_I chosen from the complex numbers. Can this restriction be removed so that λ_I is chosen from a finite set with finite entropy? In this case, a spin 1/2 particle could be completely specified with a finite amount of information. Is $C_{H,peak}$ strictly larger than $C_{H,avg}$? Another generalization includes the considerations of higher-order spin, which requires all probability moments to agree. Also, can the teleportation of continuous quantum parameters as shown in [13] be modeled by a finite-bit model. Does a general model emerge?

ACKNOWLEDGEMENTS

The author thanks Louis Sica for many valuable insights. Thanks very much also for the suggestions from Richard Cleve and Alain Tapp, and others for discussions regarding this and future work.

References

- [1] L. Vaidman, Teleporation: Dream or reality, 1998, quant-ph/9810089.
- [2] G. Brassard, R. Cleve, and A. Tapp, The cost of exactly simulating quantum entanglement with classical communication, 1999, quant-ph/9901035.
- [3] P. K. Aravind, Physics Letters A **241**, 207 (1998).
- [4] C. M. Caves and C. A. Fuchs, Quantum information: How much information in a state vector, 1996, quant-ph/9601025.
- [5] R. Hughes, *The structure and interpretation of quantum mechanics* (Harvard University Press, New York, 1989).
- [6] J. S. Bell, *Speakable and unspeakable in quantum mechanics* (Cambridge University Press, New York, 1993).

- [7] L. Devroye, *Non-Uniform Random Variate Generation* (Springer-Verlag, New York, 1986).
- [8] M. Feldmann, Foundations of Physics Letters **8**, 41 (1935).
- [9] L. Hardy, PRL **68**, 2981 (1992).
- [10] I. Percival, Quantum transfer functions, weak nonlocality and relativity, 1998, quant-ph/9803044.
- [11] A. Suarez, Relativistic nonlocality in experiments with successive impacts, 1998, quant-ph/9712049.
- [12] P. Pearle, Collapse models, 1999, quant-ph/9901077.
- [13] S. L. Braunstein and J. J. Kimble, Physical Review Letters **80**, 869 (1998).




Fig. 1: Polarization Experiment with Entangled Bell Pair and Hidden Non-Accessible Information